



National University of Technology, Islamabad
Assignment II (Calculus II), Spring 2019
Solution Key

Q1.. $\vec{AB} = -2\mathbf{i} + \mathbf{j} + \mathbf{k}$, $\vec{CD} = \mathbf{i} + 4\mathbf{j} - \mathbf{k}$, and $\vec{AC} = 2\mathbf{i} + \mathbf{j} \Rightarrow \mathbf{n} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -2 & 1 & 1 \\ 1 & 4 & -1 \end{vmatrix} = -5\mathbf{i} - \mathbf{j} - 9\mathbf{k} \Rightarrow$ the distance is

$$d = \left| \frac{(2\mathbf{i} + \mathbf{j}) \cdot (-5\mathbf{i} - \mathbf{j} - 9\mathbf{k})}{\sqrt{25 + 1 + 81}} \right| = \frac{11}{\sqrt{107}}$$

Q2.. $x = 2t$, $y = -t$, $z = -t$ represents a line containing the origin and perpendicular to the plane $2x - y - z = 4$; this line intersects the plane $3x - 5y + 2z = 6$ when t is the solution of $3(2t) - 5(-t) + 2(-t) = 6$
 $\Rightarrow t = \frac{2}{3} \Rightarrow \left(\frac{4}{3}, -\frac{2}{3}, -\frac{2}{3}\right)$ is the point of intersection

Q3. The direction of the intersection is $\mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & -2 \\ 5 & -2 & -1 \end{vmatrix} = -6\mathbf{i} - 9\mathbf{j} - 12\mathbf{k} = -3(2\mathbf{i} + 3\mathbf{j} + 4\mathbf{k})$ and is the same as the direction of the given line.

Q4. (a) The corresponding normals are $\mathbf{n}_1 = 3\mathbf{i} + 6\mathbf{k}$ and $\mathbf{n}_2 = 2\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ and since $\mathbf{n}_1 \cdot \mathbf{n}_2 = (3)(2) + (0)(2) + (6)(-1) = 6 + 0 - 6 = 0$, we have that the planes are orthogonal

(b) The line of intersection is parallel to $\mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & 0 & 6 \\ 2 & 2 & -1 \end{vmatrix} = -12\mathbf{i} + 15\mathbf{j} + 6\mathbf{k}$. Now to find a point in

the intersection, solve $\begin{cases} 3x + 6z = 1 \\ 2x + 2y - z = 3 \end{cases} \Rightarrow \begin{cases} 3x + 6z = 1 \\ 12x + 12y - 6z = 18 \end{cases} \Rightarrow 15x + 12y = 19 \Rightarrow x = 0$ and $y = \frac{19}{12}$
 $\Rightarrow \left(0, \frac{19}{12}, \frac{1}{6}\right)$ is a point on the line we seek. Therefore, the line is $x = -12t$, $y = \frac{19}{12} + 15t$ and $z = \frac{1}{6} + 6t$.

Q5.. $\mathbf{v} = (-3 \sin t)\mathbf{j} + (2 \cos t)\mathbf{k}$ and $\mathbf{a} = (-3 \cos t)\mathbf{j} - (2 \sin t)\mathbf{k}$; $|\mathbf{v}|^2 = 9 \sin^2 t + 4 \cos^2 t \Rightarrow \frac{d}{dt} (|\mathbf{v}|^2) = 18 \sin t \cos t - 8 \cos t \sin t = 10 \sin t \cos t$; $\frac{d}{dt} (|\mathbf{v}|^2) = 0 \Rightarrow 10 \sin t \cos t = 0 \Rightarrow \sin t = 0$ or $\cos t = 0$
 $\Rightarrow t = 0, \pi$ or $t = \frac{\pi}{2}, \frac{3\pi}{2}$. When $t = 0, \pi$, $|\mathbf{v}|^2 = 4 \Rightarrow |\mathbf{v}| = \sqrt{4} = 2$; when $t = \frac{\pi}{2}, \frac{3\pi}{2}$, $|\mathbf{v}| = \sqrt{9} = 3$.
Therefore max $|\mathbf{v}|$ is 3 when $t = \frac{\pi}{2}, \frac{3\pi}{2}$, and min $|\mathbf{v}| = 2$ when $t = 0, \pi$. Next, $|\mathbf{a}|^2 = 9 \cos^2 t + 4 \sin^2 t$
 $\Rightarrow \frac{d}{dt} (|\mathbf{a}|^2) = -18 \cos t \sin t + 8 \sin t \cos t = -10 \sin t \cos t$; $\frac{d}{dt} (|\mathbf{a}|^2) = 0 \Rightarrow -10 \sin t \cos t = 0 \Rightarrow \sin t = 0$ or $\cos t = 0$
 $\Rightarrow t = 0, \pi$ or $t = \frac{\pi}{2}, \frac{3\pi}{2}$. When $t = 0, \pi$, $|\mathbf{a}|^2 = 9 \Rightarrow |\mathbf{a}| = 3$; when $t = \frac{\pi}{2}, \frac{3\pi}{2}$, $|\mathbf{a}|^2 = 4 \Rightarrow |\mathbf{a}| = 2$.
Therefore, max $|\mathbf{a}| = 3$ when $t = 0, \pi$, and min $|\mathbf{a}| = 2$ when $t = \frac{\pi}{2}, \frac{3\pi}{2}$.

Q6. Let $P(t_0)$ denote the point. Then $\mathbf{v} = (5 \cos t)\mathbf{i} - (5 \sin t)\mathbf{j} + 12\mathbf{k}$ and $26\pi = \int_0^{t_0} \sqrt{25 \cos^2 t + 25 \sin^2 t + 144} dt = \int_0^{t_0} 13 dt = 13t_0 \Rightarrow t_0 = 2\pi$, and the point is $P(2\pi) = (5 \sin 2\pi, 5 \cos 2\pi, 24\pi) = (0, 5, 24\pi)$

Q7.. $\mathbf{r} = (\sqrt{2}t)\mathbf{i} + (\sqrt{2}t)\mathbf{j} + (1 - t^2)\mathbf{k} \Rightarrow \mathbf{v} = \sqrt{2}\mathbf{i} + \sqrt{2}\mathbf{j} - 2t\mathbf{k} \Rightarrow |\mathbf{v}| = \sqrt{(\sqrt{2})^2 + (\sqrt{2})^2 + (-2t)^2} = \sqrt{4 + 4t^2} = 2\sqrt{1 + t^2} \Rightarrow$ Length $= \int_0^1 2\sqrt{1 + t^2} dt = \left[2 \left(\frac{t}{2} \sqrt{1 + t^2} + \frac{1}{2} \ln \left(t + \sqrt{1 + t^2} \right) \right) \right]_0^1 = \sqrt{2} + \ln(1 + \sqrt{2})$

Q8. $\mathbf{r} = t\mathbf{i} + \ln(\cos t)\mathbf{j} \Rightarrow \mathbf{v} = \mathbf{i} + \left(\frac{-\sin t}{\cos t}\right)\mathbf{j} = \mathbf{i} - (\tan t)\mathbf{j} \Rightarrow |\mathbf{v}| = \sqrt{1^2 + (-\tan t)^2} = \sqrt{\sec^2 t} = |\sec t| = \sec t$, since $-\frac{\pi}{2} < t < \frac{\pi}{2} \Rightarrow \mathbf{T} = \frac{\mathbf{v}}{|\mathbf{v}|} = \left(\frac{1}{\sec t}\right)\mathbf{i} - \left(\frac{\tan t}{\sec t}\right)\mathbf{j} = (\cos t)\mathbf{i} - (\sin t)\mathbf{j}$; $\frac{d\mathbf{T}}{dt} = (-\sin t)\mathbf{i} - (\cos t)\mathbf{j}$
 $\Rightarrow \left| \frac{d\mathbf{T}}{dt} \right| = \sqrt{(-\sin t)^2 + (-\cos t)^2} = 1 \Rightarrow \mathbf{N} = \frac{\left(\frac{d\mathbf{T}}{dt}\right)}{\left|\frac{d\mathbf{T}}{dt}\right|} = (-\sin t)\mathbf{i} - (\cos t)\mathbf{j}$;
 $\kappa = \frac{1}{|\mathbf{v}|} \cdot \left| \frac{d\mathbf{T}}{dt} \right| = \frac{1}{\sec t} \cdot 1 = \cos t$.